Vascular Unit - VTE Risk Score

Patient details



Venous thromboembolism (VTE also known as blood clots) can be a complication of treatment. This self assessment is designed to work out whether or not you are at particularly high risk of VTE.

Medical history

Please tick every box relevant to yourself (the patient). If you are unsure about any of the questions, please ask a member of staff.

Risk			score	Risk		score	
	Age 60 year	s or above	2		Thrombophilia	3	
	Very overweight.i.e women with a waist measurement of more than				Varicose veins with phlebitis	3	
					Heart disease	1	
		and men with a	2		Lung disease	1	
	waist measurement of more than 102cm (40in)				Inflammatory disease (bowel or joints)	1	
Current medication - are you taking:				Undergoing treatment for cancer	3		
Risk			score		Previous history of leg vein clots	3	
	Oestrogen-containing contraceptive pill		2		(deep vein thrombosis) Previous history of lung clots	3	
	Hormone replacement therapy		2	_	(pulmonary embolus)	Ü	
	Tamoxifen or Erythropoietin / Darbepoetin		2		Pregnant or within 6 weeks of childbirth	3	
	Thalidomide / Lenalidomide		2		Total score:		
Fami	ily history				Total 3001c.		
Risk				score			
Onco.	A family hist (deep vein the clots (pulmon close family father, mother	2	or of	otoff in the Vaccular Unit			
Once completed, please give this form to a member of staff in the Vascular Unit. Score Recommendation							
		en active a	nd c	Irink plenty of water every day. Give the	ne		
patient the information		•					
3 or ı	As above, doctor to decide if patient needs dalteparin (Fragmin) 5,000 units once a day (as an injection) for 5 days.						
To be completed by Vascular Unit staff: scor							
	Expected significant post treatment reduction in mo				mobility?	2	
_	Procedure with a total anaesthetic and surgery time of > 90 mins or 60 mins if surgery involves pelvis or lower limbs?						
Clinical decision Total score:							
_ — — — — — — — ¬ Date:							
Stick patient label here Stick patient label here Version 3.2 Aug 2013. Approved by Thrombosis Committee.			 - -	Th Pr	o prophylaxis required. Frombotic risk outweighs bleeding risk escribe LMWH Alteparin prescribed and given		
- Thomas Committee.					(31911)		