



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



British
Thoracic
Society



COLLEGE OF
paramedics
leading the development of the paramedic profession

OXYGEN ALERT CARD



OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.

OXYGEN ALERT CARD

Name _____

I am at risk of type II respiratory failure with a raised CO₂ level. Please use my _____ % Venturi mask to achieve an oxygen saturation of _____% to _____% during exacerbations.

Use compressed air to drive nebulisers (with nasal oxygen at 2 l/min). If compressed air not available, limit oxygen-driven nebulisers to 6 minutes.